

Properties of the first $1/2^+$ state in ${}^9\text{Be}$ from electron scattering and astrophysical implications *

O. Burda, P. von Neumann-Cosel, A. Richter


Institut für Kernphysik, TU Darmstadt



TECHNISCHE
UNIVERSITÄT
DARMSTADT

- Motivation
- Experiments
- Results
- Summary and outlook

SFB 634



* Supported by the DFG within SFB 634




- ${}^9\text{Be}$ is a loosely-bound nuclear system consisting of 2α and a neutron
- ${}^9\text{Be}$ has the lowest neutron threshold ($S_n = 1.665 \text{ MeV}$) of all stable nuclei
 - first excited state lies some tens of keV above S_n
 - all excited states are unstable with respect to neutron decay

Possible role of ${}^9\text{Be}$ in the production of ${}^{12}\text{C}$



TECHNISCHE
UNIVERSITÄT
DARMSTADT




- In ν -rich environment (core-collapse supernovae) this reaction path may provide an alternative route for building up the heavy elements and triggering the ν -process

$J^\pi = 1/2^+$ state at threshold



TECHNISCHE
UNIVERSITÄT
DARMSTADT



- The photodisintegration cross section at low energies is given by the properties of $1/2^+$ resonance
- Strongly asymmetric line shape

Parameters of the first $J^\pi = 1/2^+$ state in ${}^9\text{Be}$



TECHNISCHE
UNIVERSITÄT
DARMSTADT

| | (γ, n) [1] | (e,e') [2] | [3] | Reanalysis of [2] by Barker [4] |
|------------------------------------|------------------------|---------------|----------|------------------------------------|
| E_R , MeV | 1.75(1) | 1.684(7) | 1.68(15) | 1.7316 |
| Γ_R , keV | 283(42) | 217(10) | 200(20) | 280 |
| $B(E1)\uparrow$, $e^2\text{fm}^2$ | 0.0535(35) | 0.027(2) | 0.034(3) | 0.0685 |


- [1] H. Utsunomiya et al., Phys. Rev. C63 (2001) 018801
- [2] G. Küchler et al., Z. Phys. A326 (1987) 447
- [3] J. P. Glickman et al., Phys. Rev. C43 (1991) 1740
- [4] F. C. Barker, Aust. J. Phys. 53 (2000) 247

- Resonance parameters from different experiments and analyses are not in mutual agreement

Deconvolution of the spectrum





TECHNISCHE
UNIVERSITÄT
DARMSTADT



${}^9\text{Be}(\gamma, \text{n})$ extracted from ${}^9\text{Be}(e, e')$



TECHNISCHE
UNIVERSITÄT
DARMSTADT




$$E_R = 1.737(10) \text{ MeV}$$
$$\Gamma_R = 275(14) \text{ keV}$$

Comparison: ${}^9\text{Be}(\gamma, \text{n})$ and ${}^9\text{Be}(\text{e}, \text{e}'')$



TECHNISCHE
UNIVERSITÄT
DARMSTADT



H. Utsunomiya et al.,
PRC 63, 018801

$$E_R = 1.750(10) \text{ MeV}$$
$$\Gamma_R = 283(42) \text{ keV}$$

Present


$$E_R = 1.737(10) \text{ MeV}$$
$$\Gamma_R = 275(14) \text{ keV}$$

● Resonance parameters agree well


Reaction rate $N_A \langle \sigma v \rangle$ of $\alpha + \alpha \rightarrow {}^8\text{Be}(n,\gamma){}^9\text{Be}$



TECHNISCHE
UNIVERSITÄT
DARMSTADT



Reaction rate $N_A \langle \sigma v \rangle$ of $\alpha + \alpha \rightarrow {}^8\text{Be}(n,\gamma){}^9\text{Be}$




- For $T_9 = 0.04 - 3$ the $1/2^+$ resonance determines exclusively $\alpha + \alpha \rightarrow {}^8\text{Be}(n,\gamma){}^9\text{Be}$ chain

Reaction rate $N_A \langle \sigma v \rangle$ of $\alpha + \alpha \rightarrow {}^8\text{Be}(n,\gamma){}^9\text{Be}$




TECHNISCHE
UNIVERSITÄT
DARMSTADT



Form factor of the $J^\pi = 1/2^+$ state



TECHNISCHE
UNIVERSITÄT
DARMSTADT



- NCSM: correct q dependence but difference in magnitude compared to the data

B(E1,k) strength



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Siegert's theorem:

$$B(E1,q) = (k/q)^2 B(C1,q)$$

at photon point ($q = k = E_x/\hbar c$) $B(E1,k) = B(C1,k)$

(e,e⁻) $B(C1,k) = 0.027(3) e^2 fm^2$

(γ,n) $B(E1,k) = 0.0535(35) e^2 fm^2$

- B(C1) B(E1) at photon point $k = q \rightarrow$ violation of Siegert's theorem ?

- Final values of line shape parameters of $J^\pi = 1/2^+$ resonance:
 $E_R = 1.748(6)$ MeV and $\Gamma_R = 274(8)$ keV
- $B(C1) - B(E1)$ at photon point $k = q$
→ violation of Siegert's theorem ?
- Role of direct capture
- Structure of the first $1/2^+$ state in ${}^9\text{Be}$
- NCSM and FMD calculations